

Geochemistry of radioactive elements in bituminous sands and sandstones of Permian bitumen deposits of Tatarstan (east of the Russian plate)

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 Institute of Physics Publishing. All rights reserved. The article investigates geochemical features of Permian (Cisuralian, Ufimian Stage and Biarmian, Kazanian Stage of the General Stratigraphic Scale of Russia) bituminous sands and sandstones located on the territory of the Volga-Ural oil and gas province (Republic of Tatarstan). Natural bitumens are extracted using thermal methods as deposits of high-viscosity oils. In the samples studied, the specific activity of natural radionuclides from the ^{238}U (^{226}Ra), ^{232}Th , and ^{40}K series was measured using gamma spectrometry. As a result of the precipitation of uranium and thorium and their subsequent decay, the accumulation of radium (^{226}Ra and ^{228}Ra) has been shown to occur in the bituminous substance. In the process of exploitation of bitumen-bearing rock deposits (as an oil fields) radium in the composition of a water-oil mixture can be extracted to the surface or deposited on sulfate barriers, while being concentrated on the walls of pipes and other equipment. This process requires increased attention to monitoring and inspection the environmental safety of the exploitation procedure.

<http://dx.doi.org/10.1088/1755-1315/155/1/012024>

Keywords

bitumen sands, bitumen sandstones, concentration, radioactive Elements, tar sands

References

- [1] Troyepolskiy V.I. and Lebedev N.P. 1982 Produktivnyye bituminoznyye tolshchi permskikh otlozheniy Melekesskoy vpadiny i Tatarskogo svoda (Kazan': Izdatel'stvo Kazanskogo universiteta) 104
- [2] Uspenskiy B.V. and Valeyeva I.F. 2008 Geologiya mestorozhdeniy prirodnkh bitumov Respubliki Tatarstan (Kazan': Izdatel'stvo OOO "PF "Gart") 348
- [3] Khisamov R.S., Gatiyatullin N.S. and Shargorodskiy I.Ye. (ed) 2007 Geologiya i osvoyeniye zalezhey prirodnkh bitumov Respubliki Tatarstan (Kazan': izd-vo "Fen" AN RT) 295 i dr.
- [4] Ibragimov N.G., Sizov B.A., Zajtsev, Khisamutdinov A.F. and Kubarev P.N. 2005 Radiation safety of the personnel and environmental contamination prevention at oil recovery and treating Neftyanoe Khozyaystvo - Oil Industry 12-14
- [5] Bulka G.R., Nizamutdinov N.M., Mukhutdinova N.G. et al 1991 EPR Probes in Sedimentary Rocks: The Features of Mn^{2+} and Free Radicals Distribution in the Permian Formation in Tartarstan Appl. Magn. Reson. 2 107-115
- [6] Ikeya M. 1993 World Scientific (Singapore) New Applications of Electron Spin Resonance- Dating, Dosimetry and Microscopy 509

- [7] Khasanov R.R., Mullakaev A.I. and Dusmanov E.N. 2017 The structure of sandstones in productive horizons of the Permian bituminous deposits of Tatarstan (Russia) *Uchenye Zapiski Kazanskogo Universiteta. Seriya Estestvennyye Nauki* 159 164-173 (in Russian)
- [8] Mullakaev A.I., Khasanov R.R. and Galiullin B.M. 2017 Mineralogy of sandstones and localization of oil matter in productive horizons of high-viscosity oil in permian deposits of the Volga-Ural region (Russia) 17th International Multidisciplinary Scientific GeoConference SGEM 2017, SGEM2017 Conference Proceedings 17 353-358 29 June -5 July
- [9] Conard J. 1984 *Magnetic Resonance. Introduction, Advanced Topics and Application to Fossil Energy* ed L. Petrakis and J.P. Fraissard (Dordrecht: Reidel) EPR in fossil carbonaceous materials 441-459
- [10] Jezierski A., Czechowski F., Jerzykiewicz M. and Drozd J. 2000 EPR Investigations of Structure of Humic Acids from Compost, Soil, Peat and Soft Brown Coal upon Oxidation and Metal Uptake *Appl. Magn. Reson.* 1 127-136
- [11] Mullakaev A.I., Khasanov R.R., Badrutdinov O.R., Khasanova N.M. and Nizamutdnov N.M. 2017 Features of bitumen-containing sandstones of the Volga-Ural oil and gas province according to Electron paramagnetic resonance and Gamma spectrometry data 17th International Multidisciplinary Scientific GeoConference SGEM 2017, SGEM2017 Conference Proceedings 17 133-139 29 June - 5 July
- [12] Sakhibgareev R.S. 1989 *Secondary Changes in Reservoirs during the Formation and Destruction of Oil Deposits* Leningrad, Nedra 260 (in Russian)
- [13] Zanin Y.N., Zamirailova A.G. and Eder V.G. 2016 Uranium, thorium, and potassium in black shales of the Bazhenov Formation of the West Siberian marine basin *Lithology and Mineral Resources* 51 74-85